# HPLC Application

# Tetracyclines in Meat by LC/MS/MS

Column: Dimensions: Order No: Elution Type: Eluent A: Eluent B:	50 x 4.6 mm 00B-4435-E0 Gradient		Column 50 x	4.6 mm, Ea	3				
Gradient	Step No.	Time (min)	Pct A	Pct B					
Profile:	1	0	98	2					
	2	2	98	2			Products		
	3	2.1	85	15			Froduces		
	4	6.5	50	50					
	5	7.1	0	100					
	6	8	0	100					
	7	8.1	98	2					
	8	10	98	2					
Flow Rate:	0.8 mL/min								
Col. Temp.:	40 °C	- (11 11)	(						
Detection:	Tandem Mass Spec (MS-MS) @ (600 °C)								
Detector Info:	5200 Q.1.0.0 20,								
Analyst Note:	SecurityGuard™ Guard Cartridge System extends column lifetime. - SecurityGuard Cartridges, Gemini C18 4 x 3.0mm ID, 10/Pk Part No.: AJ0-7597								
	- Holder Part No	o.: KJ0-4282							
20032				3 <					
5000 -					4				
4500 -					/				
4000 -									
3500 -					5				
3000 -			1						
2500 -			$\wedge$						
2000 -			2						
1500 -		(	$\Lambda$						
1000 -					ЛЦ				
500 -	-	m		Joh					
0 - 3	3.1 3.2 3.	3 3.4 3.5 3.6 3	.7 3.8 3.9	4 4.1 4.2	2 4.3 4.4 4.5	4.6 4.7	4.8 4.9 min		



Gemini

Products used in this application:



# **ANALYTES:**

- Oxytetracycline 1 Retention Time: 3.64 min
- 2 Tetracycline Retention Time: 3.71 min
- 3 Chlortetracycline Retention Time: 4.26 min
- Methacycline 4 Retention Time: 4.3 min
- Doxycycline 5 Retention Time: 4.42 min

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com.cn

For more information contact your Phenomenex Representative at info@phenomenex.com

for HPLC Application ID No.: 20032

# Tetracyclines in Meat by LC/MS/MS

### **PRODUCT DESCRIPTION:**

Strata<sup>™</sup>-X 33 µm Polymeric Reversed Phase, 60 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S100-UBJ

# SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

**Note:** The solvent volumes shown below are for a 60 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

#### **Condition:**

# Load:

Measure 5 g of sample into an 85 mL cenrifuge tube. Add 25 mL of acidified methanol/1M HCl mixture (0.833 mL of 1M HCl in a total volume of 200 mL methanol). Mix for 1 min using an Ultra-Turrax mixer at a velocity setting of 2. Agitate further for 5 min

#### Wash:

#### Dry:

Evaporate to dryness using an eppendorf concentrator set at 45C or blow dry with nitrogen.

#### Elute:

A

# Final Prep and Analysis:

Inject: 50 µL on HPLC Tandem Mass Spec (MS-MS) @ (600°C)

ANALYTES:		Spiked Conc. (ng/mL)	Log P	рКа	% Rec	%RSC (n=0)
1	Oxytetracycline	0				
2	Tetracycline	0				
3	Chlortetracycline	0				
4	Methacycline	0				
5	Doxycycline	0				

**Note:** This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

Ph

Phenomenex products are available worldwide.

www.phenomenex.com.cn

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at info@phenomenex.com

