HPLC Application ID No.: **23096**



Bupropion related compounds on Kinetex 5u EVO C18 150x4.6mm

Kinetex® 5µm EVO C18 100 Å, LC Column 150 x 4.6 mm, Ea

Dimensions: 150 x 4.6 mm ID Order No: 00F-4633-E0 **Elution Type:** Gradient

Eluent A: 5mM Ammonium Bicarbonate pH 9.0

Eluent B: Methanol


Gradient Step No. Time (min) Pct A Pct B **Profile:** 0 50 50 2 10 20 80

Flow Rate: 1 mL/min ambient Col. Temp.:

Detection:



Products used in this application:



ANALYTES:

Hydroxybuproprion

Retention Time: 9.156 min

2 Erythrobuproprion

Retention Time: 10.469 min

Threobupropion

Retention Time: 11.087 min

Bupropion

Retention Time: 13.939 min

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at info@phenomenex.com



www.phenomenex.com.cn info@phenomenex.com

Sample Preparation Details

for **HPLC** Application ID No.: **23096**



Bupropion related compounds on Kinetex 5u EVO C18 150x4.6mm

PRODUCT DESCRIPTION:

Strata™-X-CW 33 µm Polymeric Weak Cation, 60 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S035-UBJ

SOLID PHASE EXTRACTION	(SPE)	PRODCEDURE:
------------------------	-------	-------------

Note: The solvent volumes shown below are for a 60 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:	
Load:	
Wash:	
Dry:	
5 minutes	
Elute:	
Final Prep and Analysis:	

Inject: 2 µL on HPLC UV-Vis Abs.-Variable Wave.(UV) @ 254 nm (ambient)

ANALYTES:		Spiked Conc.	Log P	рКа	% Rec	%RSC
		(ng/mL)				(n=0)
1	Hydroxybuproprion	0				
2	Erythrobuproprion	0				
3	Threobupropion	0				
4	Bupropion	0				

This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at info@phenomenex.com

